

*Contains Nonbinding Recommendations*

**Draft Guidance on Acetazolamide**

This draft guidance, once finalized, will represent the Food and Drug Administration's (FDA's) current thinking on this topic. It does not create or confer any rights for or on any person and does not operate to bind FDA or the public. You can use an alternative approach if the approach satisfies the requirements of the applicable statutes and regulations. If you want to discuss an alternative approach, contact the Office of Generic Drugs.

**Active ingredient:** Acetazolamide

**Form/Route:** Extended Release Capsule /Oral

**Recommended studies:** 2 studies

1. Type of study: Fasting  
Design: Single-dose, two-way, crossover *in-vivo*  
Strength: 500 mg  
Subjects: Healthy males and nonpregnant females, general population.  
Additional Comments:
- 

2. Type of study: Fed  
Design: Single-dose, two-way crossover *in-vivo*  
Strength: 500 mg  
Subjects: Healthy males and nonpregnant females, general population.  
Additional Comments:
- 

**Analytes to measure (in appropriate biological fluid):** Acetazolamide in plasma

**Bioequivalence based on (90% CI):** Acetazolamide

**Waiver request of *in-vivo* testing:** Not Applicable

**Dissolution test method and sampling times:**

Please note that a **Dissolution Methods Database** is available to the public at the OGD website at <http://www.accessdata.fda.gov/scripts/cder/dissolution/>. Please find the dissolution information for this product at this website. Please conduct comparative dissolution testing on 12 dosage units each of all strengths of the test and reference products.

For modified release products, dissolution profiles generated using USP Apparatus I at 100 rpm and/or Apparatus II at 50 rpm in at least three dissolution media (pH 1.2, 4.5 and 6.8 buffers, water) should be submitted in the application. Agitation speeds may have to be increased if appropriate. It is acceptable to add a small amount of surfactant, if necessary. The following sampling times are recommended: 1, 2, and 4 hours and every 2 hours thereafter, until at least 80% of the drug is dissolved. Comparative dissolution profiles should include individual tablet data as well as the mean, range, and standard deviation at each time point for twelve tablets. Specifications will be determined upon review of the data submitted in the application.